

University of Groningen

## Physical Activity and Functional Recovery in Late-life Depression

Wassink-Vossen, Sanne

DOI:  
[10.33612/diss.147538340](https://doi.org/10.33612/diss.147538340)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2020

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*  
Wassink-Vossen, S. (2020). *Physical Activity and Functional Recovery in Late-life Depression*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.147538340>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## CHAPTER 8

### **SUMMARY AND GENERAL DISCUSSION**

This thesis studies the associations between physical activity, daily functioning and depression in older people. This knowledge is important for the development of nurse-led interventions aimed at improving functional recovery in late-life depression. To this end, four studies were conducted on a clinical cohort of depressed older subjects participating in the Netherlands Study of Depression in Older persons (NESDO). A total of 378 depressed older patients ( $\geq 60$  years of age) and a comparison group of 132 never-depressed older persons participated in NESDO.<sup>57</sup> This thesis includes up to two years of prospective data to study the course of late-life depression. In addition, to study the impact of age, we combined data from the NESDO participants with a total of 1701 depressed younger patients (18-65 years of age) who participated in the Netherlands Study of Depression and Anxiety (NESDA) using a similar methodology.<sup>58</sup> The thesis concludes with a systematic review that provides an overview of the impact of antidepressant interventions on functional recovery in late-life depression.

This chapter begins with a brief summary of the main findings of the individual studies. The first three studies, (part 1 of the thesis), focus on the relationship between physical activity and depression, while the last three studies, (part 2 of the thesis), focus on functional recovery from depression. These findings are then integrated into a series of overall conclusions. These conclusions are discussed in the light of the current literature, general clinical implications and direction for further research. This is followed by a discussion of several methodological issues that are relevant to the interpretation of findings from clinical cohort studies, including specific strengths and limitations of our studies. The thesis ends with a description of the clinical implications, primarily from the perspective of the nursing profession.

## **Summary of part 1 - Physical activity and late-life depression**

In the first part of this thesis, we focus on the association between physical activity and late-life depression. The level of physical activity can be regarded as modifiable factor that is vital for improving physical health conditions and functioning, as well as for reducing depressive symptoms in late-life (see chapter 1). An understanding of the longitudinal relation between physical activity and late life depression is important for the development of lifestyle interventions, but the determinants of physical activity in depressed older patients are still relatively unexplored. Based on cross-sectional analyses, **chapter 2** shows that depressed older patients were less physically active in comparison with their non-depressed counterparts. Adjusted for potential confounders, including demographic, health, cognitive and lifestyle characteristics, this finding remained statistically significant. In the subgroup of depressed older patients, a lower level of physical activity was independently associated with increased functional limitations, a higher number of prescribed drugs, and with inpatient treatment.

**Chapter 3**, explores the predictive contribution of physical activity and sedentary behavior on the depression outcomes at two-year follow-up. Physical activity (total MET/min per week) and sedentary behavior (total sitting min/day) were examined as determinants of late-life depression in multiple regression analyses with models adjusted for cognitive functioning, somatic conditions, psychosocial determinants, and other lifestyle characteristics. In contrast to our hypothesis, we found that in depressed older patients, neither a low level of physical activity nor sedentary behavior was associated with non-remission of the depressive disorder at two-year follow-up or with symptomatic improvement. **Chapter 4** provides possible nuances to these results. Chapter 4 examines whether age moderates the reciprocal relationship between physical activity and the course of depressive disorders over two years in patients with depressive disorders. In this study, including a sample of 1079 participants aged 18-93 years, we dichotomized the level of physical activity according to the Dutch guidelines for sufficient physical activity. Fully-adjusted regression models showed that depressed patients with sufficient physical activity at baseline were more frequently in remission at the two-year follow-up compared to depressed patients with insufficient physical activity. This main effect moderated with age. In other words, the association between sufficient physical activity and remission of the depressive disorder at two years decreases with as age increases. Another interesting finding of this study was that the level of physical activity was fairly stable over time; 74 % of the participants had not changed their level of physical activity at follow-up. Only 12 % of the participants switched from insufficient physical activity at baseline to sufficient physical activity at two-year follow-up and 14 % vice-versa. Multinomial logistic regression analyses showed that none of the potential determinants of physical activity (functional limitation, sense of mastery (chapter 3) predicted a change in physical activity over time, irrespective of age and depression outcome (i.e. diagnosis at two-year no/yes or change in IDS score after two years). The relatively low level of physical activity found in our depressed participants may therefore reflect a trait instead of a state characteristic.

## Summary of part 2 - Functional limitations and late-life depression

The second part of this thesis focuses on the relation between functional recovery and late-life depression. Both depression and increasing age are associated with functional decline. Functional disability is increasingly recognized as an important outcome measure in studies on depression, but again, knowledge on the trajectories of functioning in late-life depression is scarce. More insight into the determinants of functional recovery over time could therefore provide new insights into the needs in the treatment and care for older persons with depression. In the NESDO, functional status was measured every six months using the World Health Organization Disability Assessment Schedule 2.0 (WHODAS 2.0) self-report. The WHODAS 2.0 is a commonly used interdisciplinary

instrument that contains 36 items, covering six domains of functioning during the last 30 days: 1. Cognition (understanding and communicating); 2. Mobility (moving and getting around); 3. Hygiene, dressing, eating and staying alone (self-care); 4. Interpersonal actions (getting along with people); 5. Work, leisure, domestic responsibilities (household activities); 6. Joining in community activities (participation in society). The items regarding work were omitted for the NESDO study, because most of the participants were retired.

As shown in **Chapter 5**, the 378 depressed older patients had significantly higher levels of functional impairment compared to the 132 never-depressed older persons, as measured with WHODAS 2.0. Based on the semi-annual assessment of functional limitations, we showed that the level of functional limitations remained stable among non-remitted depressed patients and was only slightly improved among remitted depressed patients at a two-year follow-up. Moreover, remitted patients still had far more functional limitations compared to the never-depressed comparison group. The association between depression and functional impairment may suggest that it is important to include a standard clinical assessment of functioning in older subjects with depression, in line with recommendation in the DSM-5 to include measurement of the WHODAS during follow-up of anti-depressant treatment. **Chapter 6** covers the growth mixture modelling that was carried out to explore the existence of different course types of functional limitations among depressed older patients. These analyses yielded two course types, both starting at a high level of functional limitations. The smaller subgroup, consisting of only 1 in 5 depressed older patients, showed a course of functional recovery at the two-year follow-up. The largest subgroup, consisting of 4 out of 5 depressed patients, showed no functional improvement over time. The main predictor of functional recovery was symptomatic remission of depression. Combining these course types with symptomatic remission revealed four groups: 1. a subgroup with symptomatic remission and functional recovery (14 %, most optimal outcome), 2. a group of non-remitted patients without functional recovery (38 %, worst outcome), and two in-between groups consisting of either 3. patients with a symptomatic remission but no functional recovery (31 %) or 4. patients who achieved functional recovery while not being in symptomatic remission (4 %). The remaining participants (13%) were lost to follow up.

Therefore, only 3 out of 10 remitted patients also achieve functional recovery. Interestingly, 1 out of 10 patients with no symptomatic remission did achieve functional recovery. The sample size of this latter group was too small for multivariable analyses, but the finding implies that functional recovery is possible even in the presence of an ongoing, chronic depressive disorder.

Predictors of lack of functional recovery were then explored by multinomial logistic regression, stratified by symptomatic remission yes/no. We found that patients who remained depressed and showed no functional recovery were characterized by the

presence of more chronic somatic diseases, a lower sense of mastery, and higher levels of anxiety, when compared to patients with the most optimal results (reference group). Among symptomatic remitted patients, the lack of functional recovery was associated with female gender, higher educational level, higher gait speed, and lower depressive symptom severity.

Since only 20% of depressed older patients overall were found to achieve functional recovery over a two-year period, we argue that the outcome of mental health interventions should include functional recovery in addition to symptomatic remission.

**Chapter 7** therefore covers that systematic review that was carried out to evaluate the effectiveness of interventions on functional limitations in patients with late-life depression. The 14 randomized controlled trials that were identified by the literature search evaluated four different treatment strategies, i.e. psychological interventions ( $n=7$ ), pharmacotherapy ( $n=3$ ), physical exercise ( $n=3$ ), and collaborative care ( $N=4$ ). Whereas all of these treatment strategies showed improvement with respect to depressive symptoms, collaborative-care interventions had the most favorable effect on the reduction of functional limitations in late-life depression. To optimize anti-depressant treatment by improving functional recovery, collaborative or integrated care should be the core when treating late-life depression.

## Main findings of the thesis

Since the primary objectives of this thesis are discussed extensively in the separate chapters, this section discusses the four main conclusions based on the different studies. First, we found that the effect of sufficient physical activity on the course of late-life depression remains ambiguous. Second, physical activity seems to reflect a trait rather than a state in late-life depression (and probably at all ages). Third, symptomatic remission of depression does not imply functional recovery in late-life depression. This leads to a fourth conclusion, namely that integrated care is needed for full recovery from late-life depression, and future research should focus on recovery of daily functioning instead of the severity of depressive symptoms only. These four main conclusions will be discussed in more detail below.

### The impact of physical activity on the course of late-life depression: enough is enough?

Sufficient physical activity is associated with improved physical health as well as mental health. Likewise, this thesis confirmed that depressed older patients are less physically active than their non-depressed counterparts (chapter 2). Sufficient physical activity can thus be seen as a protective factor for not developing depressive disorder.<sup>248,249</sup> In this

section, we will argue that sufficient physical activity may also be seen as a predictive factor for remission of depression in clinically depressed patients of all ages. We also discuss that the impact of physical activity on the course of late-life depression remains ambiguous regarding the contradictory results in chapter 3 and 4.

While in chapter 3 no effect of physical activity on late-life depression outcome could be demonstrated, in chapter 4, insufficient physical activity was found to predict an adverse outcome of depression in both younger and older patients. However, this effect was more pronounced in the younger compared to the older age group (chapter 4). The most likely explanation for this contradictory finding in late-life depression is the fact that physical activity was operationalized in two different ways, i.e. total amount of physical activity in MET-minutes in chapter 3 and a dichotomization of physical activity according to the WHO-definition of “sufficient” physical activity.<sup>77</sup> It is important to note that sufficient physical activity is not simply the application of a cut-off on the number of MET-minutes, because the relative contribution of light, moderate and heavy physical activity also has to be taken into account. To examine the discrepant findings in chapter 3 and 4, in chapter 4, we conducted post-hoc sensitivity analyses with the total amount of physical activity (as in chapter 3). These sensitivity analyses showed that a higher total MET-minutes/week at baseline increased the odds of remission at follow up, but in line with chapter 3, this effect was lost in the fully adjusted model and was not moderated by age.

Apparently, the predictive effect of physical activity on remission of depression is characterized by a subthreshold - ‘enough is enough’ - and not a dose-response relation. Once the level of sufficient physical activity is reached, additional activity does not have an additional effect on depression. This suggests that increasing the level of physical activity from insufficient to sufficient might be helpful in the treatment of depression. This finding is in line with studies in younger depressed patients, in which it was demonstrated that the protective effect of additional exercises is found among persons with relatively low levels of physical activity, with no indication of any additional benefit beyond a certain amount of physical activity a week.<sup>250,251</sup>

The absence of any effect of physical activity on depression outcomes in chapter 3 may also be explained by a lack of statistical power. The strength of this association decreases with age, which implies that a higher number of subjects is needed to demonstrate a statistically significant effect. Nevertheless, patient numbers were markedly lower in the higher age-ranges. Lastly, this age effect may also be explained by the fact that older patients with depression have more pre-existing chronic diseases<sup>165</sup> and the additional high level of functional limitations improves only minimally after remission of the depressive disorder in older people (chapter 5). A course with functional recovery was found in only one fifth of the older patients (chapter 6). The presence of these functional limitations in later life could also explain why improving physical activity can be more difficult in later life.

In conclusion, older subjects with late-life depression have mostly adapted to a lifestyle with a low activity level that is hard to change, partly due to physical inabilities. It is likely that insufficient physical activity predisposes them to developing of depression, and when depressed, sufficient physical activity could lead to a favorable course of depression.

## **physical activity in late-life depression reflects a trait rather than a state characteristic**

Physical activity is regarded as a key factor with respect to modifiable behavior to improve physical and mental health. Nevertheless, this section argues that physical activity primarily reflects a trait characteristic of depressed patients rather than a state-effect due to the depressive disorder. This statement unfortunately implies that physical activity behavior will be hard to change in late-life depression.

Contrary to our expectations, the level of physical activity hardly changed over time (chapter 4) and this was independent of depression outcome. Neither remission of the depressive disorder, nor a decline in depressive symptoms over time predicted a change in physical activity at follow-up. In the case of a state-effect of the depressive disorder on the level of physical activity, one would expect a dose-response effect. However, the severity of the depression was not associated with the level of physical activity (chapter 2). These findings suggest that the lower level of physical activity found among depressed patients compared to their never-depressed counterparts merely reflects a trait characteristic. This interpretation offers an explanation for the fact that although lifestyle interventions are feasible and may look attractive because they target potentially modifiable behavior, results until now have been rather meager. Physical exercise interventions thus far have only demonstrated marginal short-term effects in the treatment of depression in adults and older adults.<sup>252,253</sup> Thus far, a long-term effect of exercise interventions on depression has not been found.<sup>41</sup> It is important to note that, adherence has been identified as an explanatory factor for the heterogeneity of results of the effect of exercise programs on depressive symptoms by meta-regression analysis of RCT's in late life depression.<sup>252</sup> In other words, high adherence to exercise is a prerequisite for the effectiveness of such lifestyle interventions in reducing depressive symptoms.<sup>254</sup> If the level of physical activity indeed reflects merely a trait rather than a state characteristic, it remains questionable to what extent older adults (with depressed mood) are capable of long-term, sustainable behavioral change with high adherence.



## **Symptomatic recovery does not reflect functional recovery in late life depression**

The burden of psychiatric disorders is more dependent on the level of limitations in daily activities and relevant social contacts than on the actual severity of the psychiatric symptoms.<sup>166</sup> Functional recovery should therefore be an additional treatment target along with symptomatic recovery. We have studied the longitudinal course of functioning in late-life depression and searched for determinants of functional improvement over time. By combining symptomatic and functional recovery in the analyses, we showed that symptomatic recovery of late -life depression does not parallel functional recovery. Functional limitations improve only minimally after the remission of depressive symptoms (chapter 5), and functional recovery is only achieved by about one fifth of depressed older patients (chapter 6).

We do not know to what extent pre-existing functional limitations, caused by age-related factors, contribute to this unfavorable course. We assume that depression itself is a contributor to this outcome, because functional deficits also often remain in younger, somatically healthy depressed adults, even among patients who achieve symptomatic remission.<sup>168</sup> However, among younger depressed patients functional improvement is more directly associated with a reduction in depressive symptoms.<sup>255</sup> This suggests that depression itself causes persistent functional impairment in subgroups, irrespective of age.

Two important determinants of functional recovery are relevant in geriatric psychiatry: depression itself (also relevant for younger patients as described above) and somatic morbidity. We found that a higher number of chronic diseases was associated with non-recovery of functional limitation (see chapter 5 and 6). This is likely to be more specific for old age, because in old age, comorbidity with somatic diseases is more prominent than at younger ages.<sup>165</sup> In the NESDO study, a higher number of comorbid somatic diseases was associated with a poor course of late-life depression.<sup>2,256</sup> Moreover, a quarter of the sample could be classified as physically frail<sup>225</sup> which was also associated with a detrimental course of the depressive disorder.<sup>169</sup> Depression moderates the level of functional impairment associated with many somatic diseases, including cardiac disease, morbus Parkinson, and mild cognitive impairment.<sup>188,257,258</sup> These findings would suggest that improving depression will also improve the level of functional impairment, albeit not to the level of healthy older persons (as found in chapter 5). However, the chronicity of depressive symptoms in the face of multimorbidity and frailty may also point to over-detection of depression, as it has been argued that depressive symptoms are simply a sign or epiphenomenon of these underlying conditions.<sup>259</sup>

In conclusion, we found that functional status remains impaired in most of our subjects at the two year follow-up and that this is associated with more somatic illnesses. On

the other hand, we also found a group that was rather robust but nevertheless severely depressed and functionally impaired at baseline. Interestingly, this group showed the most favorable course in functional as well as symptomatic recovery (chapter 6). For future studies it would be interesting to separate these two groups; the old, frail with high levels of functional limitations versus the younger and fit/robust, with severe depressive disorders.

Furthermore, single-intervention anti-depressant treatment is apparently not enough to achieve functional recovery for a large group of patients. For optimal functional recovery, integrated care that includes the treatment of multiple comorbid medical and social problems simultaneously is needed. This type of treatment aligns with the multifactorial origin of (functional) impairment.

## **Towards integrated care with a focus on functioning**

The multifactorial origin of impairment in the older and frail depressed population stresses the need for a view that is broader than just the disease-centered approach. A focus on health and functioning as key outcomes seems to be more appropriate. Only 14 studies have examined which interventions may be relevant to improve functional recovery in late-life depression (Chapter 7). The evidence points towards collaborative care programs in particular, to increase the level of functioning of older people with depression.

Collaborative care studies were conducted in the early 1990s in the USA.<sup>260</sup> These models of integrated care are designed to promote collaboration between primary care providers and mental health specialists to improve access, planning, quality, and outcomes and treatment of depressed older adults. The studies included in our review were effective in improving functional limitations, and case management conducted mainly by specialized nurses was a common feature. Unfortunately, we could not determine which treatment components precisely predict functional improvement in these studies. In our opinion, however, it may not be specific interventions, but the personalized integrated care plan. Future studies into collaborative care should conduct a 'complex-intervention-studies approach' in order to disentangle the effects of the components of the intervention.<sup>261</sup>

The PRISM-e study showed that for major depression, reduction in symptom severity was higher for those randomly assigned to the enhanced specialty referral group in comparison with integrated care in primary care, whereas change in functioning did not differ across the two care models.<sup>262</sup> This may suggest that for major depression, treatment in specialized care is still indicated (according to psychiatric guidelines), but it needs more personalized care management within integrated care to improve functioning in collaboration with primary care. Translating this to the current Dutch organization of healthcare, where investments have been made in primary care to enable depression

treatment, in the next step, a focus on the collaboration of mental health institutions and primary care is needed.


This integrated approach to treat depressive disorder by incorporating strategies to simultaneously target comorbid somatic diseases, functional impairment and taking personal preferences and strength and weaknesses into account, is nothing new.<sup>263</sup> The paradigm shift from a biomedical model towards a more inclusive definition of health has been accomplished in the past decades and resulted in various theoretical concepts like the ICF,<sup>4</sup> the “positive health” definition<sup>202</sup> and the positive psychology trend in mental health care.<sup>264,265</sup>

The concept of the International Classification of Functioning, Disability and Health (ICF) was reported to be useful in guiding clinical assessment and providing a comprehensive understanding of health and functioning from the first-person perspective. It considers a wide range of domains of functioning, which is particularly relevant for complex conditions.<sup>125</sup> Another proposal that takes a broader look at health-outcomes, is the “dynamic concept of health” by Huber et al.<sup>202</sup> She defines health as “the ability to adapt and to self-manage in the light of social, physical, and emotional challenges”. This broad perception, which is captured in six domains (bodily functions, mental functions and perception, spiritual dimension, quality of life, social and societal participation and daily function) still requires further operationalization for clinical practice. It could result in a new clinical practice where patients are involved in making shared decisions based on the different health dimensions. This would result in improvements in the ‘ability to cope’, ‘resilience’ and ‘self-management’, as well as in better physical and mental health, in functioning in the various dimensions, and/or in health-related quality of life.

Since depressed older patients are a highly heterogeneous group with respect to underlying risk factors, comorbidities and consequences, integrated personalized care will optimize daily functioning (Chapter 7). In our study, patients with neither symptomatic remission nor functional recovery were characterized by a higher somatic disease burden, anxiety, and low sense of mastery. For these patients, geriatric rehabilitation should be considered as one of the core components of integrated care. And, according to our results in chapter 2 and 6, from a psychosocial perspective, it might be relevant to incorporate interventions with focus on a sense of mastery and anxiety. This approach provides the opportunity to add to the currently available anti-depressant treatment interventions, and to improve the success rate of anti-depressant therapy. By extending psychiatric symptom counts as a treatment outcome with functional outcomes, treatment outcomes are measured in a way that is closed to the patient’s definition of a successful treatment.

Nevertheless, integrated care is not yet mainstream in everyday practice and scientific research. Organizational, financial, and technical hurdles need to be overcome for a solid reorganization from disease management towards integrated network health-care.<sup>266</sup>

Partnerships across traditional organizational and professional boundaries are needed. In an integrated system these partnerships can pass through the boundaries of the ‘cure’ and ‘care’ sector to provide a real continuum of care.<sup>267</sup> In the case of depression in older adults, this involves a reassessment of the role of specialized mental healthcare. Instead of delivering treatment from isolated outpatient psychiatric clinics, the system moves towards following the individual patient with his/her individual needs, in partnership with primary care and other care providers, in a network around the patient. In this scenario, mental health nurses could play an important role in the maintenance of symptomatic and functional recovery and by serving as case-managers between the mental health system and the other members in the network of the individual patient to optimize functional recovery.



**Figure 1:** Relations found in this thesis

## Methodological considerations

The empirical findings in this thesis are based primarily on data from the Netherlands Study of Depression in Older persons (NESDO). In chapter 4, this data was augmented with data on depressed younger patients participating in the Netherlands Study of Depression and Anxiety (NESDA). Clinical cohort studies have several strengths and limitations. We will discuss the following aspects of these studies, which should be taken into account when weighing the outcome of this thesis: the definition of depression, the use of self-report questionnaires, especially in the assessment of physical activity and functional impairment, and finally the black box of interventions patients have received.

### Case definition of depression

The main subject of our studies was depression in later life. NESDO provides a large sample of older persons with depression. To include a broad range of depression severity, subjects were recruited from in primary care as well as (in- and outpatient) specialized mental health care. After recruitment, the depression diagnosis was confirmed by trained research assistants using the CIDI. The 378 subjects included depressed participants in the NESDO study experiencing a current DSM-IV classification of minor depression (5.6%) or major depressive disorder (95%), of which 26.5 % presented comorbid dysthymia. In this thesis, all analyzed samples included all depression diagnoses. In both samples (NESDA and NESDO), current psychiatric diagnosis was defined by a recency of 6 months. In the cross-sectional study (chapter 2), we selected only depressed cases who also met the criteria for a depressive disorder in the past month, while the level of activity was assessed over the previous week. This led to a more valid comparison of physical activity (since recent remission may already have increased the level of physical activity in previous week), but also to a smaller sample due to the exclusion of 72 participants. Based on the results in chapter 4, however, including all patients would not have biased the results, because physical activity appeared to be merely a trait characteristic. In the chapters that use longitudinal data, the importance of recency was considered to be less important than the opportunity to include as many as possible participants with follow up data. Moreover, by including all depressed patients using the past 6-month definition, we followed the a-priori definition of depression of NESDO and NESDA. Since physical activity appeared to remain stable over time and change in physical activity was not associated with remission of the depressive disorder at two-year follow-up, it is unlikely that these differences in recency affected the results.

One of the strong points of our dataset is that, in addition to the formal depression diagnosis, we also had measures on depression symptom severity, such as the IDS scores. The results on these scores show that we did include participants with a broad range of severity of depressive symptoms,<sup>57</sup> which reflects clinical practice. We were also

able to analyze the bidirectional relations between various depression characteristics in relation with physical activity and functional limitations.

## Outcome measurements

One of the strengths of observational cohort studies, especially NESDO and NESDA, is the opportunity to measure a plethora of data. This enabled us to study determinants of physical activity and functional limitations in late-life depression from a biopsychosocial perspective over time. We were nevertheless bound to the questionnaires used in NESDO. Physical activity was measured with the International Physical Activity Questionnaire (IPAQ) and functional limitations with the WHODAS 2.0, both self-report questionnaires. This section will cover usability and the cons regarding the use of our two major outcome measurements.

**Physical activity** – Physical activity was measured with the IPAQ. The IPAQ is a well-accepted instrument that has been validated in various subpopulations around the world.<sup>77</sup> Compared to other self-reported physical activity questionnaires, the IPAQ is one of the questionnaires with the most acceptable psychometric properties.<sup>105</sup> Reliability studies have shown that the IPAQ can be used in studies with repeated measurements, although the true magnitude of the change over time, if any, may not be accurate.<sup>133</sup> Results, especially in older adults, are mixed. For example, the IPAQ had poor concurrent validity and only categorized physical behavior correctly in 2% of older cases.<sup>268</sup>

Compared to the measurement of physical activity through the use of wearables, the IPAQ tends to overestimate the amount of physical activity in (healthy) older people.<sup>269</sup> Kowalski et al.<sup>270</sup> found only a moderate correlation between objective and self-reported physical activity in older adults ( $r=0.38$ ). These results, however, are difficult to extrapolate to older persons with depression, because negative cognitions could downgrade this perception and cognitive problems could preclude a valid judgement even more.<sup>271</sup> Schuch et al.<sup>272</sup> found that the total time spent on physical activity was significantly higher with the use of objective measurements than with self-reported measurements in adults with depression. This suggests an underestimation of physical activity in depression, in contrast to the overestimation found among non-depressed persons. In our study, however, self-reported physical activity hardly changed over two years in an initially depressed group, and this was not influenced by remission of depression or by age (Chapter 4). In line with our self-report measurements, objectively measured physical activity studies also showed that depressed patients are less physically active in comparison to their non-depressed counterparts.<sup>114</sup>

In conclusion, it is evident that self-reported physical activity questionnaires measure the perception of one's own activity instead of the actual amount of activity. However,

wearables were not in widespread use at the start of NESDO. The increasingly available and affordable accelerometry devices would therefore be a logical addition in future research.<sup>273,274</sup> The results on the difference in perception (questionnaire) and the actual amount of activity (accelerometry) would therefore provide valuable information to boost (or develop) motivational interventions for lifestyle change as part of treatment.

**Functional limitations** - A recent systematic review on the assessment of the instrumental activities of daily living (IADL), social functioning, and general functioning showed that the large majority of instruments have not been validated in patients suffering from late-life depression.<sup>240,242</sup> Based on their findings, these researchers developed an algorithm to select functional assessment instruments in late-life depression for researchers and clinicians. With regard to the assessment of general functioning, the WHODAS 2.0 and the Late-Life Function and Disability Instrument (LLFDI) were recommended for research and clinical use. The LLFDI has more robust evidence for this subgroup, but was considered the second choice because the LLFDI requires more training and expertise to use. The LLFDI could therefore be used for researchers assessing functioning in late-life depression as a primary outcome, whereas the WHODAS 2.0 is the best alternative as a secondary outcome or supplemental part of their study.

Worldwide the WHODAS 2.0 is well-known and used in research and daily practice across all age groups.<sup>275</sup> In 2014, the American Psychiatric Association added the WHODAS 2.0 to the DSM-5 clinical classification to replace the General Assessment of Functioning (GAF) score. Despite the fact that work-related items were omitted for the NESDO study in order to optimize the sensitivity for change in functioning in retired older adults, results showed little change in WHODAS 2.0 total scores over time (chapter 5 and 6). To detect subtle changes, more research is needed in analyzing subscales, as was done in chapter 5. Unfortunately, due to skewed distribution of subscale scores (high frequency of null scores), this was not possible in the GMM analysis in chapter 6.

## **Longitudinal observational study**

The strength of this thesis is the opportunity to use the longitudinal data of the NESDO and NESDA. This allowed us to look at changes over time in chapters 3-6 and draw cause and consequence conclusions. The NESDO and NESDA were observational studies in which no restrictions were imposed on regular clinical care. Since the majority of patients were recruited after referral to specialized mental health care, most subjects have definitely received some form of treatment. Although the use of medication (including antidepressants and other psychotropic drugs) was available at baseline and two-year follow-up, it was not possible to assess the effect and appropriateness of these and other interventions on the course of the depressive disorder. First of all, at baseline it was not known whether patients had previously received anti-depressant

medication for a prolonged period (e.g. referred for treatment resistance) or were recently switched to these drugs. Furthermore, no data were available on the duration of use, dosage regimes and switches during the two-years of follow-up. Information on the other interventions, especially psychotherapy or interventions aimed at improving lifestyle, such as food and activity-habits, would be very interesting, but were not available for our groups, so we could only speculate as to the effect of these interventions. Nonetheless, in the Netherlands we have evidence-based guidelines for the treatment of depression, including late-life depression, and empirical studies have shown that adherence to these guidelines is quite good.<sup>276</sup> If true in our sample, we must conclude that the overall results of these interventions on the clinical course of depression were rather disappointing. However, the actual nature of these interventions remains a 'black box', as is the case in many observational studies similar to ours.

The meager results of treatment and the predominantly poor outcome after two and six years in our NESDO group could reflect the inherent problems of old-age psychiatry.<sup>165</sup> Most depressed subjects had somatic co-morbidity. In addition to depression, a range of other, somatic and cognitive/degenerative problems are involved, all with a negative impact on outcomes. This is also reflected in the rather large numbers of drop-outs at follow up, due to death (6.8% after 2 years, 16.4% after 6 years) and inability to continue participation (9.7% after 2 years, 11,5% after 6 years).<sup>2,16</sup> Participants lost to follow up were more severely depressed at baseline and had more functional limitations than the participants that were included in the analytical samples, suggesting that the most severely impaired persons were not followed-up. This may have biased our results towards an underestimation of the association that was found between depression and functional limitations. Furthermore, our study did not assess pre-morbid levels of functioning. We therefore cannot take a possible pre-existing vulnerability into account.

## Clinical implications

In the previous paragraphs, much is said about the implications of our results for clinical practice and research. It is clear that treatment and care for patients with late-life depression is complex and demands efforts from multiple disciplines. Future research should aim to indicate what works for whom. The starting point of this thesis was to contribute to the body of knowledge on the interrelationship between physical activity, functional recovery, and depressive disorder. Nurses play a major role in interventions aimed at these daily activities and functional recovery. The acquired knowledge could help to develop more targeted and nurse-led interventions. Therefore, in this last section, we will focus on the implications of this thesis for clinical care and research from the perspective of the nursing professionals.



With regard to physical activity, the challenge is to nudge patients towards a long term, sustainable behavioral change with high adherence. Based on this thesis, we can assume that enough is enough (chapters 3 and 4). Patients with low daily physical activity, can therefore be motivated to increase their level to a personalized amount of sufficient physical activity. Integrating physical activity into daily routines is emerging as a promising alternative.<sup>277</sup> This is currently being studied in the PreventIT project, which addresses the different phases of adopting a healthier lifestyle.<sup>278</sup> Based on our results, for patients suffering from late-life depression, it is important to take into account determinants of physical activity such as a low sense of mastery and individual functional impairments. Extra attention is also needed with inpatients (chapter 2). Nurses are well educated and are in a position to encourage and assist patients to engage in daily physical activity. Since the level of physical activity is merely a trait instead of a state characteristic (chapter 4), there will be a group of patients who requires long-term encouragement and support to attain this goal.


Concerning functional recovery, this thesis shows that only 2 out of 10 patients functionally recover, which can be interpreted as a large gap between symptomatic remission and functional recovery. One of the main conclusions of this thesis is therefore that increased focus on functional recovery as outcome in practice and research is inevitable. As mentioned earlier, functioning as an outcome is the key-factor for nursing professionals, who generally focus on the human as a person functioning in his or her daily life and not as someone with a specific disease. We therefore advocate, in line with Stallinga,<sup>279</sup> the adoption, practice and research of functioning as core outcome for the nursing profession. The ICF, with its acknowledgement of the interaction between people and their environments in health and disability, is a useful conceptual framework for clinical reasoning of nurses.<sup>280-282</sup>

A specific finding in this thesis, was that the most vulnerable and functionally-impaired depressed patients are characterized by high prevalence of chronic somatic diseases, a lower sense of mastery, and a higher level of anxiety (chapter 6). This vulnerability and comorbidity may have a negative impact on the self-management skills of patients. Self-management abilities, however, are important as older persons generally face a large network of caregivers due to their multimorbidity. The opportunity for optimizing functional recovery is a multidisciplinary collaboration of patient-centered, integrated care (chapter 7).

It is conceivable that these patients will benefit from the involvement of a nurse who takes on a crucial role in assisting the patient and their families in managing and coordinating care. This means helping the patients to define their unmet needs, assist them in goal setting, guide them in the information needed for informed shared decision making in treatment options, support them in their process and evaluate the effect of interventions

together. In such a way *'that the patient would perform unaided if they had the necessary strength, will, or knowledge and to do this in such a way as to help them gain full of partial independence as rapidly as possible'*.<sup>3</sup>

Being a nurse practitioner myself, I hope this thesis will stimulate and contribute the development of integrated care for depressed older persons beyond the borders of mental and somatic health care.



## REFERENCES

1. Kok RM, Reynolds CF, 3rd. Management of depression in older adults: A review. *JAMA*. 2017;317(20):2114-2122.
2. Comijs HC, Nieuwesteeg J, Kok R, et al. The two-year course of late-life depression; results from the Netherlands study of depression in older persons. *BMC Psychiatry*. 2015;15:20-015-0401-5.
3. Henderson V. The nature of nursing. *Am J Nurs*. 1964;64:62-68.
4. World Health Organization. International classification of functioning, disability and health. . 2001.
5. Bize R, Johnson JA, Plotnikoff RC. Physical activity level and health-related quality of life in the general adult population: A systematic review. *Prev Med*. 2007;45(6):401-415.
6. World Health Organization. Global recommendations on physical activity for health, 65 years and above. <https://www.who.int/dietphysicalactivity/physical-activity-recommendations-65years.pdf?ua=1>). Updated 2011.
7. Trimbos-instituut. Addendum ouderen bij de Multidisciplinaire richtlijn depressie. 2011.
8. NICE. *Depression: The treatment and management of depression in adults (updated edition)*. Leicester UK: The British Psychological Society & The Royal College of Psychiatrists, 2010; 2010. <http://dx.doi.org/>.
9. Nguyen HQ, Koepsell T, Unutzer J, Larson E, LoGerfo JP. Depression and use of a health plan-sponsored physical activity program by older adults. *Am J Prev Med*. 2008;35(2):111-117.
10. Mitnitski AB, Mogilner AJ, Rockwood K. Accumulation of deficits as a proxy measure of aging. *ScientificWorldJournal*. 2001;1:323-336.
11. Wang L, van Belle G, Kukull WB, Larson EB. Predictors of functional change: A longitudinal study of nondemented people aged 65 and older. *J Am Geriatr Soc*. 2002;50(9):1525-1534.
12. Barry LC, Allore HG, Bruce ML, Gill TM. Longitudinal association between depressive symptoms and disability burden among older persons. *J Gerontol A Biol Sci Med Sci*. 2009;64(12):1325-1332.
13. Lenze EJ, Schulz R, Martire LM, et al. The course of functional decline in older people with persistently elevated depressive symptoms: Longitudinal findings from the cardiovascular health study. *J Am Geriatr Soc*. 2005;53(4):569-575.
14. Lenze EJ, Rogers JC, Martire LM, et al. The association of late-life depression and anxiety with physical disability: A review of the literature and prospectus for future research. *Am J Geriatr Psychiatry*. 2001;9(2):113-135.
15. World Health Organization. Factsheet depression. <https://www.who.int/en/news-room/factsheets/detail/depression>. Updated 2020.
16. Jeuring HW, Stek ML, Huisman M, et al. A six-year prospective study of the prognosis and predictors in patients with late-life depression. *Am J Geriatr Psychiatry*. 2018;26(9):985-997.
17. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 5th ed. Washington, DC: Author; 2013.
18. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 4th ed. Washington, DC: Author; 2000.
19. Krishnan KR, Hays JC, Tupler LA, George LK, Blazer DG. Clinical and phenomenological comparisons of late-onset and early-onset depression. *Am J Psychiatry*. 1995;152(5):785-788.

20. Corruble E, Gorwood P, Falissard B. Association between age of onset and symptom profiles of late-life depression. *Acta Psychiatr Scand*. 2008;118(5):389-394.
21. Alexopoulos GS. Depression in the elderly. *Lancet*. 2005;365(9475):1961-1970.
22. Naarding P, Beekman AT. Vascular depression: Where do we go from here? *Expert Rev Neurother*. 2011;11(1):77-83.
23. Alexopoulos GS, Kelly RE, Jr. Research advances in geriatric depression. *World Psychiatry*. 2009;8(3):140-149.
24. Bauman A, Schoeppe S, Lewicka M. Review of best practice in interventions to promote physical activity in developing countries; Background document prepared for the WHO workshop on physical activity and public health. . 2008.
25. Miller ME, Rejeski WJ, Reboussin BA, Ten Have TR, Ettinger WH. Physical activity, functional limitations, and disability in older adults. *J Am Geriatr Soc*. 2000;48(10):1264-1272.
26. Hall KS, Morey MC, Dutta C, et al. Activity-related energy expenditure in older adults: A call for more research. *Med Sci Sports Exerc*. 2014;46(12):2335-2340.
27. Owen N. Sedentary behavior: Understanding and influencing adults' prolonged sitting time. *Prev Med*. 2012;55(6):535-539.
28. Teychenne M, Ball K, Salmon J. Sedentary behavior and depression among adults: A review. *Int J Behav Med*. 2010;17(4):246-254.
29. Wirth K, Klenk J, Brefka S, et al. Biomarkers associated with sedentary behaviour in older adults: A systematic review. *Ageing Res Rev*. 2017;35:87-111.
30. Chastin SF, Buck C, Freiburger E, et al. Systematic literature review of determinants of sedentary behaviour in older adults: A DEDIPAC study. *Int J Behav Nutr Phys Act*. 2015;12:127-015-0292-3.
31. Ainsworth BE. How do I measure physical activity in my patients? questionnaires and objective methods. *Br J Sports Med*. 2009;43(1):6-9.
32. Molarius A, Berglund K, Eriksson C, et al. Mental health symptoms in relation to socio-economic conditions and lifestyle factors--a population-based study in sweden. *BMC Public Health*. 2009;9:302-2458-9-302.
33. Pisinger C, Toft U, Aadahl M, Glumer C, Jorgensen T. The relationship between lifestyle and self-reported health in a general population: The Inter99 study. *Prev Med*. 2009;49(5):418-423.
34. Suetani S, Stubbs B, McGrath JJ, Scott JG. Physical activity of people with mental disorders compared to the general population: A systematic review of longitudinal cohort studies. *Soc Psychiatry Psychiatr Epidemiol*. 2019;54(12):1443-1457.
35. Schuch FB, Vancampfort D, Firth J, et al. Physical activity and incident depression: A meta-analysis of prospective cohort studies. *Am J Psychiatry*. 2018;175(7):631-648.
36. de Rezende LF, Rodrigues Lopes M, Rey-Lopez JP, Matsudo VK, Luiz Odo C. Sedentary behavior and health outcomes: An overview of systematic reviews. *PLoS One*. 2014;9(8):e105620.
37. Stubbs B, Vancampfort D, Hallgren M, et al. EPA guidance on physical activity as a treatment for severe mental illness: A meta-review of the evidence and position statement from the european psychiatric association (EPA), supported by the international organization of physical therapists in mental health (IOPTMH). *Eur Psychiatry*. 2018;54:124-144.

38. Sarris J, O'Neil A, Coulson CE, Schweitzer I, Berk M. Lifestyle medicine for depression. *BMC Psychiatry*. 2014;14:107-244X-14-107.
39. Cooney GM, Dwan K, Greig CA, et al. Exercise for depression. *Cochrane Database Syst Rev*. 2013;(9):CD004366. doi(9):CD004366.
40. van Gool CH, Kempen GI, Penninx BW, Deeg DJ, Beekman AT, van Eijk JT. Relationship between changes in depressive symptoms and unhealthy lifestyles in late middle aged and older persons: Results from the longitudinal aging study amsterdam. *Age Ageing*. 2003;32(1):81-87.
41. Krogh J, Nordentoft M, Sterne JA, Lawlor DA. The effect of exercise in clinically depressed adults: Systematic review and meta-analysis of randomized controlled trials. *J Clin Psychiatry*. 2011;72(4):529-538.
42. Blake H, Mo P, Malik S, Thomas S. How effective are physical activity interventions for alleviating depressive symptoms in older people? A systematic review. *Clin Rehabil*. 2009;23(10):873-887.
43. Daskalopoulou C, Stubbs B, Kralj C, Koukounari A, Prince M, Prina AM. Physical activity and healthy ageing: A systematic review and meta-analysis of longitudinal cohort studies. *Ageing Res Rev*. 2017;38:6-17.
44. McAuley E, Szabo AN, Mailey EL, et al. Non-exercise estimated cardiorespiratory fitness: Associations with brain structure, cognition, and memory complaints in older adults. *Ment Health Phys Act*. 2011;4(1):5-11.
45. Pahor M, Guralnik JM, Ambrosius WT, et al. Effect of structured physical activity on prevention of major mobility disability in older adults: The LIFE study randomized clinical trial. *JAMA*. 2014;311(23):2387-2396.
46. Reiner M, Niermann C, Jekauc D, Woll A. Long-term health benefits of physical activity--a systematic review of longitudinal studies. *BMC Public Health*. 2013;13:813-2458-13-813.
47. Win S, Parakh K, Eze-Nliam CM, Gottdiener JS, Kop WJ, Ziegelstein RC. Depressive symptoms, physical inactivity and risk of cardiovascular mortality in older adults: The cardiovascular health study. *Heart*. 2011;97(6):500-505.
48. Iliffe S, Kharicha K, Carmaciu C, et al. The relationship between pain intensity and severity and depression in older people: Exploratory study. *BMC Fam Pract*. 2009;10:54-2296-10-54.
49. Karsten J, Hartman CA, Ormel J, Nolen WA, Penninx BW. Subthreshold depression based on functional impairment better defined by symptom severity than by number of DSM-IV symptoms. *J Affect Disord*. 2010;123(1-3):230-237.
50. Samson MM, Meeuwse IB, Crowe A, Dessens JA, Duursma SA, Verhaar HJ. Relationships between physical performance measures, age, height and body weight in healthy adults. *Age Ageing*. 2000;29(3):235-242.
51. van der Voort TY, Seldenrijk A, van Meijel B, et al. Functional versus syndromal recovery in patients with major depressive disorder and bipolar disorder. *J Clin Psychiatry*. 2015;76(6):e809-14.
52. Verhaak PF, Dekker JH, de Waal MW, van Marwijk HW, Comijs HC. Depression, disability and somatic diseases among elderly. *J Affect Disord*. 2014;167:187-191.
53. McKnight PE, Kashdan TB. The importance of functional impairment to mental health outcomes: A case for reassessing our goals in depression treatment research. *Clinical Psychology Review*. 2009(29):243-259.

54. ten Doesschate MC, Koeter MW, Bockting CL, Schene AH, DELTA Study Group. Health related quality of life in recurrent depression: A comparison with a general population sample. *J Affect Disord.* 2010;120(1-3):126-132.
55. Geerlings SW, Beekman AT, Deeg DJ, Twisk JW, Van Tilburg W. The longitudinal effect of depression on functional limitations and disability in older adults: An eight-wave prospective community-based study. *Psychol Med.* 2001;31(8):1361-1371.
56. World Health Organization. How to use the ICF: A practical manual for using the international classification of functioning, disability and health (ICF). exposure draft for comment. . 2013.
57. Comijs HC, van Marwijk HW, van der Mast RC, et al. The netherlands study of depression in older persons (NESDO); a prospective cohort study. *BMC Res Notes.* 2011;4:524-0500-4-524.
58. Penninx BW, Beekman AT, Smit JH, et al. The netherlands study of depression and anxiety (NESDA): Rationale, objectives and methods. *Int J Methods Psychiatr Res.* 2008;17(3):121-140.
59. World Health Organization. Global strategy on diet, physical activity and health. <http://www.who.int/dietphysicalactivity/pa/en/index.html>. Updated 2011. Accessed februari 3, 2014.
60. Booth FW, Chakravarthy MV, Gordon SE, Spangenburg EE. Waging war on physical inactivity: Using modern molecular ammunition against an ancient enemy. *Journal of applied Physiology.* 2002;93(1):3-30.
61. Pruimboom L. Physical inactivity is a disease synonymous for a non-permissive brain disorder. *Medical Hypotheses.* 2011;77:708-713.
62. Benedetti TR, Borges LJ, Petroski EL, Goncalves LH. Physical activity and mental health status among elderly people. *Rev Saude Publica.* 2008;42(2):302-307.
63. Lindwall M, Larsman P, Hagberg MS. The reciprocal relationship between physical activity and depression in older european adults: A prospective cross-lagged panel design using SHARE data. *Health Psychol.* 2011;30(4):453-462.
64. Uher R, Perlis RH, Henigsberg N, et al. Depression symptom dimensions as predictors of antidepressant treatment outcome: Replicable evidence for interest-activity symptoms. *Psychol Med.* 2012;42(5):967-980.
65. Sjosten N, Kivela SL. The effects of physical exercise on depressive symptoms among the aged: A systematic review. *Int J Geriatr Psychiatry.* 2006;21(5):410-418.
66. Warburton DER, Bredin SSD. Health benefits of physical activity: A systematic review of current systematic reviews. *Curr Opin Cardiol.* 2017;32(5):541-556.
67. Koenen MA, Verheijden MW, Chinapaw MJ, Hopman-Rock M. Determinants of physical activity and exercise in healthy older adults: A systematic review. *Int J Behav Nutr Phys Act.* 2011;8:142-5868-8-142.
68. van Stralen MM, de Vries H, Mudde AN, Bolman C, Lechner L. Efficacy of two tailored interventions promoting physical activity in older adults. *Am J Prev Med.* 2009;37(5):405-417.
69. Blumenthal JA, Babyak MA, Doraiswamy PM, et al. Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med.* 2007;69(7):587-596.
70. van Stralen MM, Lechner L, Mudde AN, de Vries H, Bolman C. Determinants of awareness, initiation and maintenance of physical activity among the over-fifties: A delphi study. *Health Educ Res.* 2010;25(2):233-247.



71. Hoffman BM, Blumenthal JA, Babyak MA, et al. Exercise fails to improve neurocognition in depressed middle-aged and older adults. *Med Sci Sports Exerc.* 2008;40(7):1344-1352.
72. Antonelli Incalzi R, Corsonello A, Pedone C, Corica F, Carbonin P. Depression and drug utilization in an elderly population. *Therapeutics and Clinical Risk Management.* 2005;1(1):55-60.
73. Duivis HE, de Jonge P, Penninx BW, Na BY, Cohen BE, Whooley MA. Depressive symptoms, health behaviors, and subsequent inflammation in patients with coronary heart disease: Prospective findings from the heart and soul study. *Am J Psychiatry.* 2011;168(9):913-920.
74. Harvey SB, Hotopf M, Overland S, Mykletun A. Physical activity and common mental disorders. *Br J Psychiatry.* 2010;197(5):357-364.
75. Trost SG, Owen N, Bauman AE, Sallis JF, Brown W. Correlates of adults' participation in physical activity: Review and update. *Med Sci Sports Exerc.* 2002;34(12):1996-2001.
76. Craig CL, Marshall AL, Sjoström M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc.* 2003;35(8):1381-1395.
77. Patterson E. Guidelines for data processing and analysis of the international Physical Activity questionnaire (IPAQ). <https://sites.google.com/site/theipaq/scoring-protocol>. Updated 2005. Accessed July/19, 2017.
78. Ainsworth BE, Haskell WL, Whitt MC, et al. Compendium of physical activities: An update of activity codes and MET intensities. *Medicine & Science in Sports & Exercise.* 2000;32(9):S498-504.
79. Tomioka K, Iwamoto J, Saeki K, Okamoto N. Reliability and validity of the international physical activity questionnaire (IPAQ) in elderly adults: The fujiwara-kyo study. *J Epidemiol.* 2011;21(6):459-465.
80. Andrews G, Peters L. The psychometric properties of the composite international diagnostic interview. *Soc Psychiatry Psychiatr Epidemiol.* 1998;33(2):80-88.
81. Rush AJ, Gullion CM, Basco MR, Jarrett RB, Trivedi MH. The inventory of depressive symptomatology (IDS): Psychometric properties. *Psychol Med.* 1996;26(3):477-486.
82. Hegeman JM, Wardenaar KJ, Comijs HC, de Waal MW, Kok RM, van der Mast RC. The subscale structure of the inventory of depressive symptomatology self report (IDS-SR) in older persons. *J Psychiatr Res.* 2012;46(10):1383-1388.
83. Korten NC, Penninx BW, Kok RM, et al. Heterogeneity of late-life depression: Relationship with cognitive functioning. *Int Psychogeriatr.* 2014;26(6):953-963.
84. Klein M, Ponds RW, Houx PJ, Jolles J. Effect of test duration on age-related differences in stroop interference. *J Clin Exp Neuropsychol.* 1997;19(1):77-82.
85. Chwastiak LA, Von Korff M. Disability in depression and back pain: Evaluation of the world health organization disability assessment schedule (WHO DAS II) in a primary care setting. *J Clin Epidemiol.* 2003;56(6):507-514.
86. Kriegsman DM, Penninx BW, van Eijk JT, Boeke AJ, Deeg DJ. Self-reports and general practitioner information on the presence of chronic diseases in community dwelling elderly. A study on the accuracy of patients' self-reports and on determinants of inaccuracy. *J Clin Epidemiol.* 1996;49(12):1407-1417.
87. Babor TF, Kranzler HR, Lauerma RJ. Early detection of harmful alcohol consumption: Comparison of clinical, laboratory, and self-report screening procedures. *Addict Behav.* 1989;14(2):139-157.

88. Stansfeld S, Marmot M. Deriving a survey measure of social support: The reliability and validity of the close persons questionnaire. *Soc Sci Med*. 1992;35(8):1027-1035.
89. Pearlin LI, Schooler C. The structure of coping. *J Health Soc Behav*. 1978;19(1):2-21.
90. Cairney J, Faulkner G, Veldhuizen S, Wade TJ. Changes over time in physical activity and psychological distress among older adults. *Can J Psychiatry*. 2009;54(3):160-169.
91. Rimer J, Dwan K, Lawlor DA, et al. Exercise for depression. *Cochrane Database Syst Rev*. 2012;(7):CD004366. doi(7):CD004366.
92. Bridle C, Spanjers K, Patel S, Atherton NM, Lamb SE. Effect of exercise on depression severity in older people: Systematic review and meta-analysis of randomised controlled trials. *Br J Psychiatry*. 2012;201(3):180-185.
93. van Stralen, M. M., De Vries, H., Mudde, H. N., Bolman, C., Lechner, L. Determinants of initiation and maintenance of physical activity among older adults: A literature review. . 2009.
94. Penninx BW, Leveille S, Ferrucci L, van Eijk JT, Guralnik JM. Exploring the effect of depression on physical disability: Longitudinal evidence from the established populations for epidemiologic studies of the elderly. *Am J Public Health*. 1999;89(9):1346-1352.
95. van't Veer-Tazelaar PJ, van Marwijk HW, van Oppen P, et al. Stepped-care prevention of anxiety and depression in late life: A randomized controlled trial. *Arch Gen Psychiatry*. 2009;66(3):297-304.
96. King AC, King DK. Physical activity for an aging population. *Public health reviews*. 2010;32(2):401-426.
97. Happell B, Platania-Phung C, Scott D. Placing physical activity in mental health care: A leadership role for mental health nurses. *Int J Ment Health Nurs*. 2011;20(5):310-318.
98. Ferrucci L, Schrack JA, Knuth ND, Simonsick EM. Aging and the energetic cost of life. *J Am Geriatr Soc*. 2012;60(9):1768-1769.
99. Richardson CR, Faulkner G, McDevitt J, Skrinar GS, Hutchinson DS, Piette JD. Integrating physical activity into mental health services for persons with serious mental illness. *Psychiatr Serv*. 2005;56(3):324-331.
100. Studenski S, Carlson MC, Fillit H, Greenough WT, Kramer A, Rebok GW. From bedside to bench: Does mental and physical activity promote cognitive vitality in late life? *Sci Aging Knowledge Environ*. 2006;2006(10):pe21.
101. Underwood M, Lamb SE, Eldridge S, et al. Exercise for depression in elderly residents of care homes: A cluster-randomised controlled trial. *Lancet*. 2013;382(9886):41-49.
102. Pilling S, Anderson I, Goldberg D, Meader N, Taylor C, Two Guideline Development Groups. Depression in adults, including those with a chronic physical health problem: Summary of NICE guidance. *BMJ*. 2009;339:b4108.
103. Rosenberg D, Depp CA, Vahia IV, et al. Exergames for subsyndromal depression in older adults: A pilot study of a novel intervention. *Am J Geriatr Psychiatry*. 2010;18(3):221-226.
104. Grimm EK, Swartz AM, Hart T, Miller NE, Strath SJ. Comparison of the IPAQ-short form and accelerometry predictions of physical activity in older adults. *J Aging Phys Act*. 2012;20(1):64-79.
105. Forsen L, Loland NW, Vuillemin A, et al. Self-administered physical activity questionnaires for the elderly: A systematic review of measurement properties. *Sports Med*. 2010;40(7):601-623.

106. Hamer M, Stamatakis E. Prospective study of sedentary behavior, risk of depression, and cognitive impairment. *Med Sci Sports Exerc.* 2014;46(4):718-723.
107. Steketee G. Methodological issues in prediction of treatment outcome. *Clinical Psychology Review.* 1992;12(4):387-400.
108. Strawbridge WJ, Deleger S, Roberts RE, Kaplan GA. Physical activity reduces the risk of subsequent depression for older adults. *Am J Epidemiol.* 2002;156(4):328-334.
109. Boschloo L, Reeuwijk KG, Schoevers RA, W J H Penninx B. The impact of lifestyle factors on the 2-year course of depressive and/or anxiety disorders. *J Affect Disord.* 2014;159:73-79.
110. Harris AH, Cronkite R, Moos R. Physical activity, exercise coping, and depression in a 10-year cohort study of depressed patients. *J Affect Disord.* 2006;93(1-3):79-85.
111. Beekman AT, Copeland JR, Prince MJ. Review of community prevalence of depression in later life. *Br J Psychiatry.* 1999;174:307-311.
112. Mammen G, Faulkner G. Physical activity and the prevention of depression: A systematic review of prospective studies. *Am J Prev Med.* 2013;45(5):649-657.
113. Wassink-Vossen S, Noorthoorn EO, Collard RM, Comijs HC, Oude Voshaar RC, Naarding P. Value of physical activity and sedentary behavior in predicting depression in older adults. *J Am Geriatr Soc.* 2016;64(3):647-649.
114. Wielopolski J, Reich K, Clepce M, et al. Physical activity and energy expenditure during depressive episodes of major depression. *J Affect Disord.* 2015;174:310-316.
115. Hiles SA, Lamers F, Milaneschi Y, Penninx BWJH. Sit, step, sweat: Longitudinal associations between physical activity patterns, anxiety and depression. *Psychol Med.* 2017;47(8):1466-1477.
116. Wassink-Vossen S, Collard RM, Oude Voshaar RC, Comijs HC, de Vocht HM, Naarding P. Physical (in)activity and depression in older people. *J Affect Disord.* 2014;161:65-72.
117. Rethorst CD, Trivedi MH. Evidence-based recommendations for the prescription of exercise for major depressive disorder. *J Psychiatr Pract.* 2013;19(3):204-212.
118. Silveira H, Moraes H, Oliveira N, Coutinho ES, Laks J, Deslandes A. Physical exercise and clinically depressed patients: A systematic review and meta-analysis. *Neuropsychobiology.* 2013;67(2):61-68.
119. Chi WC, Chang KH, Escorpizo R, et al. Measuring disability and its predicting factors in a large database in taiwan using the world health organization disability assessment schedule 2.0. *Int J Environ Res Public Health.* 2014;11(12):12148-12161.
120. Sargent-Cox KA, Butterworth P, Anstey KJ. Role of physical activity in the relationship between mastery and functional health. *Gerontologist.* 2015;55(1):120-131.
121. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12(3):189-198.
122. Wittchen HU. Reliability and validity studies of the WHO--composite international diagnostic interview (CIDI): A critical review. *J Psychiatr Res.* 1994;28(1):57-84.
123. Wittchen HU, Burke JD, Semler G, Pfister H, Von Cranach M, Zaudig M. Recall and dating of psychiatric symptoms. test-retest reliability of time-related symptom questions in a standardized psychiatric interview. *Arch Gen Psychiatry.* 1989;46(5):437-443.
124. Rush AJ, Giles DE, Schlessner MA, Fulton CL, Weissenburger J, Burns C. The inventory for depressive symptomatology (IDS): Preliminary findings. *Psychiatry Res.* 1986;18(1):65-87.

125. Ustun TB, Chatterji S, Kostanjsek N, et al. Developing the world health organization disability assessment schedule 2.0. *Bull World Health Organ*. 2010;88(11):815-823.
126. de Beurs E, Comijs H, Twisk JW, Sonnenberg C, Beekman AT, Deeg D. Stability and change of emotional functioning in late life: Modelling of vulnerability profiles. *J Affect Disord*. 2005;84(1):53-62.
127. Pearlin LI, Nguyen KB, Schieman S, Milkie MA. The life-course origins of mastery among older people. *J Health Soc Behav*. 2007;48(2):164-179.
128. Thombs BD, Ziegelstein RC, Pilote L, et al. Somatic symptom overlap in beck depression inventory-II scores following myocardial infarction. *Br J Psychiatry*. 2010;197(1):61-66.
129. Hagstromer M, Kwak L, Oja P, Sjostrom M. A 6 year longitudinal study of accelerometer-measured physical activity and sedentary time in swedish adults. *J Sci Med Sport*. 2015;18(5):553-557.
130. Ku PW, Fox KR, Chen LJ, Chou P. Physical activity and depressive symptoms in older adults: 11-year follow-up. *Am J Prev Med*. 2012;42(4):355-362.
131. Franek J. Self-management support interventions for persons with chronic disease: An evidence-based analysis. *Ont Health Technol Assess Ser*. 2013;13(9):1-60.
132. Peckham AD, McHugh RK, Otto MW. A meta-analysis of the magnitude of biased attention in depression. *Depress Anxiety*. 2010;27(12):1135-1142.
133. Lee PH, Macfarlane DJ, Lam TH, Stewart SM. Validity of the international physical activity questionnaire short form (IPAQ-SF): A systematic review. *Int J Behav Nutr Phys Act*. 2011;8:115-5868-8-115.
134. Soundy A, Roskell C, Stubbs B, Vancampfort D. Selection, use and psychometric properties of physical activity measures to assess individuals with severe mental illness: A narrative synthesis. *Arch Psychiatr Nurs*. 2014;28(2):135-151.
135. Mota-Pereira J, Silverio J, Carvalho S, Ribeiro JC, Fonte D, Ramos J. Moderate exercise improves depression parameters in treatment-resistant patients with major depressive disorder. *J Psychiatr Res*. 2011;45(8):1005-1011.
136. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med*. 2006;3(11):e442.
137. Heo M, Murphy CF, Fontaine KR, Bruce ML, Alexopoulos GS. Population projection of US adults with lifetime experience of depressive disorder by age and sex from year 2005 to 2050. *Int J Geriatr Psychiatry*. 2008;23(12):1266-1270.
138. Bickenbach J, Cieza A, Rauch, A., Stucki, G. *ICF core sets; manual for clinical practice*. Göttingen, Germany: Hogrefe publishing; 2012.
139. Riddle M, Potter GG, McQuoid DR, Steffens DC, Beyer JL, Taylor WD. Longitudinal cognitive outcomes of clinical phenotypes of late-life depression. *Am J Geriatr Psychiatry*. 2017;25(10):1123-1134.
140. World Health Organization. *Composite international diagnostic interview. version 2.1*. Geneva: WHO; 1997.
141. Kessler RC, Birnbaum HG, Shahly V, et al. Age differences in the prevalence and co-morbidity of DSM-IV major depressive episodes: Results from the WHO world mental health survey initiative. *Depress Anxiety*. 2010;27(4):351-364.

142. Wittchen HU, Robins LN, Cottler LB, Sartorius N, Burke JD, Regier D. Cross-cultural feasibility, reliability and sources of variance of the composite international diagnostic interview (CIDI). the multicentre WHO/ADAMHA field trials. *Br J Psychiatry*. 1991;159:645-53, 658.
143. World Health Organization. Disability assessment schedule II (WHODAS II). *WHO*. 2000.
144. Garin O, Ayuso-Mateos JL, Almansa J, et al. Validation of the "world health organization disability assessment schedule, WHODAS-2" in patients with chronic diseases. . 2010;8:51.
145. Mackin RS, Ayalon L, Feliciano L, Arean PA. The sensitivity and specificity of cognitive screening instruments to detect cognitive impairment in older adults with severe psychiatric illness. *J Geriatr Psychiatry Neurol*. 2010;23(2):94-99.
146. Cockrell JR, Folstein MF. Mini-mental state examination (MMSE). *Psychopharmacol Bull*. 1988;24(4):689-692.
147. Cuijpers P, de Graaf R, van Dorsselaer S. Minor depression: Risk profiles, functional disability, health care use and risk of developing major depression. *J Affect Disord*. 2004;79(1-3):71-79.
148. Cuijpers P, Smit F. Subthreshold depression as a risk indicator for major depressive disorder: A systematic review of prospective studies. *Acta Psychiatr Scand*. 2004;109(5):325-331.
149. Bijl RV, Ravelli A, van Zessen G. Prevalence of psychiatric disorder in the general population: Results of the netherlands mental health survey and incidence study (NEMESIS). *Soc Psychiatry Psychiatr Epidemiol*. 1998;33(12):587-595.
150. Braam AW, Copeland JR, Delespaul PA, et al. Depression, subthreshold depression and comorbid anxiety symptoms in older europeans: Results from the EURODEP concerted action. *J Affect Disord*. 2014;155:266-272.
151. Licht-Strunk E, van der Windt DA, van Marwijk HW, de Haan M, Beekman AT. The prognosis of depression in older patients in general practice and the community. A systematic review. *Fam Pract*. 2007;24(2):168-180.
152. Stek ML, Van Exel E, Van Tilburg W, Westendorp RG, Beekman AT. The prognosis of depression in old age: Outcome six to eight years after clinical treatment. *Aging Ment Health*. 2002;6(3):282-285.
153. Barry LC, Murphy TE, Gill TM. Depressive symptoms and functional transitions over time in older persons. *Am J Geriatr Psychiatry*. 2011;19(9):783-791.
154. Ormel J, Oldehinkel AJ, Nolen WA, Vollebergh W. Psychosocial disability before, during, and after a major depressive episode: A 3-wave population-based study of state, scar, and trait effects. *Arch Gen Psychiatry*. 2004;61(4):387-392.
155. Buist-Bouwman MA, Ormel J, de Graaf R, Vollebergh WA. Functioning after a major depressive episode: Complete or incomplete recovery? *J Affect Disord*. 2004;82(3):363-371.
156. Rhebergen D, Beekman AT, de Graaf R, et al. Trajectories of recovery of social and physical functioning in major depression, dysthymic disorder and double depression: A 3-year follow-up. *J Affect Disord*. 2010;124(1-2):148-156.
157. Karppinen H, Laakkonen ML, Strandberg TE, Huohvanainen EA, Pitkala KH. Do you want to live to be 100? answers from older people. *Age Ageing*. 2016;45(4):543-549.
158. Bruce ML, Lohman MC, Greenberg RL, Bao Y, Raue PJ. Integrating depression care management into medicare home health reduces risk of 30- and 60-day hospitalization: The depression care for patients at home cluster-randomized trial. *J Am Geriatr Soc*. 2016;64(11):2196-2203.

159. Maarsingh OR, Heymans MW, Verhaak PF, Penninx BWJH, Comijs HC. Development and external validation of a prediction rule for an unfavorable course of late-life depression: A multicenter cohort study. *J Affect Disord*. 2018;235:105-113.
160. Apostolo J, Bobrowicz-Campos E, Rodrigues M, Castro I, Cardoso D. The effectiveness of non-pharmacological interventions in older adults with depressive disorders: A systematic review. *Int J Nurs Stud*. 2016;58:59-70.
161. Gilbody S, Lewis H, Adamson J, et al. Effect of collaborative care vs usual care on depressive symptoms in older adults with subthreshold depression: The CASPER randomized clinical trial. *JAMA*. 2017;317(7):728-737.
162. Luppá M, Sikorski C, Luck T, et al. Age- and gender-specific prevalence of depression in latest-life--systematic review and meta-analysis. *J Affect Disord*. 2012;136(3):212-221.
163. Kamenov K, Cabello M, Coenen M, Ayuso-Mateos JL. How much do we know about the functional effectiveness of interventions for depression? A systematic review. *J Affect Disord*. 2015;188:89-96.
164. Hoogendijk EO, Deeg DJ, Poppelaars J, et al. The longitudinal aging study amsterdam: Cohort update 2016 and major findings. *Eur J Epidemiol*. 2016;31(9):927-945.
165. Schaakxs R, Comijs HC, Lamers F, Kok RM, Beekman ATF, Penninx BWJH. Associations between age and the course of major depressive disorder: A 2-year longitudinal cohort study. *Lancet Psychiatry*. 2018;5(7):581-590.
166. Zimmerman M, McGlinchey JB, Posternak MA, Friedman M, Attiullah N, Boerescu D. How should remission from depression be defined? the depressed patient's perspective. *Am J Psychiatry*. 2006;163(1):148-150.
167. Prodingier B, Cieza A, Oberhauser C, et al. Toward the international classification of functioning, disability and health (ICF) rehabilitation set: A minimal generic set of domains for rehabilitation as a health strategy. *Arch Phys Med Rehabil*. 2016;97(6):875-884.
168. Sheehan DV, Nakagome K, Asami Y, Pappadopoulos EA, Boucher M. Restoring function in major depressive disorder: A systematic review. *J Affect Disord*. 2017;215:299-313.
169. Collard RM, Wassink-Vossen S, Schene AH, et al. Symptomatic and functional recovery in depression in later life. *Soc Psychiatry Psychiatr Epidemiol*. 2018.
170. Meltzer H, Bebbington P, Brugha T, et al. Physical ill health, disability, dependence and depression: Results from the 2007 national survey of psychiatric morbidity among adults in england. *Disabil Health J*. 2012;5(2):102-110.
171. Kessler RC, Berglund P, Demler O, et al. The epidemiology of major depressive disorder. *J.Am.Med.Dir.Assoc*. 2003;280(23).
172. Ekers D, Murphy R, Archer J, Ebenezer C, Kemp D, Gilbody S. Nurse-delivered collaborative care for depression and long-term physical conditions: A systematic review and meta-analysis. *J Affect Disord*. 2013;149(1-3):14-22.
173. Van der Elst W, van Boxtel MP, van Breukelen GJ, Jolles J. Rey's verbal learning test: Normative data for 1855 healthy participants aged 24-81 years and the influence of age, sex, education, and mode of presentation. *J Int Neuropsychol Soc*. 2005;11(3):290-302.
174. Walton D. The diagnostic and predictive accuracy of the modified word learning test in psychiatric patients over 65. *J Ment Sci*. 1958;104(437):1119-1122.

175. Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: Psychometric properties. *J Consult Clin Psychol*. 1988;56:893-897.
176. Andruff H, Carraro N, Thompson A, Gaudreau P. Latent growth modelling: A tutorial. *Tutorials in Quantitative Methods for Psychology*. 2009;5(1):11-24.
177. Muthén LK, Muthén BO. *Mplus user's guide*. Vol 7. Los Angeles CA: Muthén & Muthén; 2012.
178. Raab M, Fasang A, Hess M. Pathways to death: The co-occurrence of physical and mental health in the last years of life. *demographic research*. 2018;38(53):1619-1634.
179. Hybels CF, Pieper CF, Blazer DG, Fillenbaum GG, Steffens DC. Trajectories of mobility and IADL function in older patients diagnosed with major depression. *Int J Geriatr Psychiatry*. 2010;25(1):74-81.
180. Tunvirachaisakul C, Gould RL, Coulson MC, et al. Predictors of treatment outcome in depression in later life: A systematic review and meta-analysis. *J Affect Disord*. 2018;227:164-182.
181. Hammer-Helmich L, Haro JM, Jonsson B, et al. Functional impairment in patients with major depressive disorder: The 2-year PERFORM study. *Neuropsychiatr Dis Treat*. 2018;14:239-249.
182. Jaeger J, Berns S, Uzelac S, Davis-Conway S. Neurocognitive deficits and disability in major depressive disorder. *Psychiatry Res*. 2006;145(1):39-48.
183. Friborg O, Martinsen EW, Martinussen M, Kaiser S, Overgard KT, Rosenvinge JH. Comorbidity of personality disorders in mood disorders: A meta-analytic review of 122 studies from 1988 to 2010. *J Affect Disord*. 2014;152-154:1-11.
184. Koorevaar AM, Comijs HC, Dhondt AD, et al. Big five personality and depression diagnosis, severity and age of onset in older adults. *J Affect Disord*. 2013;151(1):178-185.
185. Zimmermann T, Puschmann E, van den Bussche H, et al. Collaborative nurse-led self-management support for primary care patients with anxiety, depressive or somatic symptoms: Cluster-randomised controlled trial (findings of the SMADS study). *Int J Nurs Stud*. 2016;63:101-111.
186. Wermelinger Avila MP, Lucchetti AL, Lucchetti G. Association between depression and resilience in older adults: A systematic review and meta-analysis. *Int J Geriatr Psychiatry*. 2017;32(3):237-246.
187. Hybels CF, Blazer DG, Landerman LR, Steffens DC. Heterogeneity in symptom profiles among older adults diagnosed with major depression. *Int Psychogeriatr*. 2011;23(6):906-922.
188. Hybels CF, Pieper CF, Blazer DG, Steffens DC. Heterogeneity in the three-year course of major depression among older adults. *Int J Geriatr Psychiatry*. 2016;31(7):775-782.
189. IsHak WW, Bonifay W, Collison K, et al. The recovery index: A novel approach to measuring recovery and predicting remission in major depressive disorder. *J Affect Disord*. 2017;208:369-374.
190. Holvast F, Massoudi B, Oude Voshaar RC, Verhaak PFM. Non-pharmacological treatment for depressed older patients in primary care: A systematic review and meta-analysis. *PLoS One*. 2017;12(9):e0184666.



191. Ferrari AJ, Charlson FJ, Norman RE, et al. Burden of depressive disorders by country, sex, age, and year: Findings from the global burden of disease study 2010. *PLoS Med*. 2013;10(11):e1001547.
192. Tedeschini E, Levkovitz Y, Iovieno N, Ameral VE, Nelson JC, Papakostas GI. Efficacy of antidepressants for late-life depression: A meta-analysis and meta-regression of placebo-controlled randomized trials. *J Clin Psychiatry*. 2011;72(12):1660-1668.
193. Calati R, Salvina Signorelli M, Balestri M, et al. Antidepressants in elderly: Metaregression of double-blind, randomized clinical trials. *J Affect Disord*. 2013;147(1-3):1-8.
194. Wassink-Vossen S, Collard RM, Wardenaar KJ, et al. Trajectories and determinants of functional limitations in late-life depression: A 2-year prospective cohort study. *Eur Psychiatry*. 2019;62:90-96.
195. Bickenbach J. Ethics, disability and the international classification of functioning, disability and health. *Am J Phys Med Rehabil*. 2012;91(13 Suppl 1):S163-7.
196. Hudelson PM. Quality research for health programmes. 1994.
197. United Nations, Department of Economic and Social Affairs, Population Division. World population ageing 2015. . 2015;ST/ESA/SER.A/390.
198. Baune BT, Adrian I, Jacobi F. Medical disorders affect health outcome and general functioning depending on comorbid major depression in the general population. *J Psychosom Res*. 2007;62(2):109-118.
199. Cronin-Stubbs D, de Leon CF, Beckett LA, Field TS, Glynn RJ, Evans DA. Six-year effect of depressive symptoms on the course of physical disability in community-living older adults. *Arch Intern Med*. 2000;160(20):3074-3080.
200. Licht-Strunk E, van der Windt DA, van Marwijk HW, de Haan M, Beekman AT. The prognosis of depression in older patients in general practice and the community. A systematic review. *Fam Pract*. 2007;24(2):168-180.
201. Lenze EJ, Miller MD, Dew MA, et al. Subjective health measures and acute treatment outcomes in geriatric depression. *Int J Geriatr Psychiatry*. 2001;16(12):1149-1155.
202. Huber M, Knottnerus JA, Green L, et al. How should we define health? *BMJ*. 2011;343:d4163.
203. European Commission. Horizon 2020. <https://ec.europa.eu/programmes/horizon2020/en>. Updated 2020.
204. Mind. Mind platform. [www.mindplatform.nl/over-mind-platform](http://www.mindplatform.nl/over-mind-platform). Updated 2020.
205. Rijksoverheid. Ministerie van volksgezondheid, welzijn en sport. <https://www.rijksoverheid.nl/ministeries/ministerie-van-volksgezondheid-welzijn-en-sport>. Updated 2020.
206. Boevink W. Tree: Towards recovery, empowerment and experiential expertise of users of psychiatric services. In: Ryan P, Ramon S, Greacen T, eds. *Empowerment, lifelong learning and recovery in mental health*. 1st ed. UK: Palgrave Macmillan; 2012.
207. Langlieb AM, Guico-Pabia CJ. Beyond symptomatic improvement: Assessing real-world outcomes in patients with major depressive disorder. *Prim Care Companion J Clin Psychiatry*. 2010;12(2):10.4088/PCC.09r00826blu.
208. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev*. 2015;4:1-4053-4-1.



209. Higgins JP, Altma, Gotzsche PC, et al. The cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928.
210. Higgins JPT, Savović j, Page PJ, Sterne JAC. Revised cochrane risk-of-bias tool for randomized trials (RoB 2). . 2019.
211. Tarlov AR, Ware JE,Jr, Greenfield S, Nelson EC, Perrin E, Zubkoff M. The medical outcomes study. an application of methods for monitoring the results of medical care. *JAMA*. 1989;262(7):925-930.
212. Alexopoulos GS, Raue P, Arean P. Problem-solving therapy versus supportive therapy in geriatric major depression with executive dysfunction. *Am J Geriatr Psychiatry*. 2003;11(1):46-52.
213. Unutzer J, Katon W, Callahan CM, et al. Collaborative care management of late-life depression in the primary care setting: A randomized controlled trial. *JAMA*. 2002;288(22):2836-2845.
214. Brenes GA, Williamson JD, Messier SP, et al. Treatment of minor depression in older adults: A pilot study comparing sertraline and exercise. *Aging Ment Health*. 2007;11(1):61-68.
215. Choi NG, Marti CN, Bruce ML, Hegel MT, Wilson NL, Kunik ME. Six-month postintervention depression and disability outcomes of in-home telehealth problem-solving therapy for depressed, low-income homebound older adults. *Depress Anxiety*. 2014;31(8):653-661.
216. Gitlin LN, Harris LF, McCoy MC, et al. A home-based intervention to reduce depressive symptoms and improve quality of life in older african americans: A randomized trial. *Ann Intern Med*. 2013;159(4):243-252.
217. Huang TT, Liu CB, Tsai YH, Chin YF, Wong CH. Physical fitness exercise versus cognitive behavior therapy on reducing the depressive symptoms among community-dwelling elderly adults: A randomized controlled trial. *Int J Nurs Stud*. 2015;52(10):1542-1552.
218. Kiosses DN, Arean PA, Teri L, Alexopoulos GS. Home-delivered problem adaptation therapy (PATH) for depressed, cognitively impaired, disabled elders: A preliminary study. *Am J Geriatr Psychiatry*. 2010;18(11):988-998.
219. van Marwijk HW, Ader H, de Haan M, Beekman A. Primary care management of major depression in patients aged > or =55 years: Outcome of a randomised clinical trial. *Br J Gen Pract*. 2008;58(555):680-6, I-II; discussion 687.
220. Williams JW, Barrett J, Oxman T, et al. Treatment of dysthymia and minor depression in primary care: A randomized controlled trial in older adults. *JAMA*. 2000;284(12):1519-1526.
221. Neviani F, Belvederi Murri M, Mussi C, et al. Physical exercise for late life depression: Effects on cognition and disability. *Int Psychogeriatr*. 2017;29(7):1105-1112.
222. Rondanelli M, Giacosa A, Opizzi A, et al. Long chain omega 3 polyunsaturated fatty acids supplementation in the treatment of elderly depression: Effects on depressive symptoms, on phospholipids fatty acids profile and on health-related quality of life. *J Nutr Health Aging*. 2011;15(1):37-44.
223. Alexopoulos GS, Raue PJ, Kiosses DN, Seirup JK, Banerjee S, Arean PA. Comparing engage with PST in late-life major depression: A preliminary report. *Am J Geriatr Psychiatry*. 2015;23(5):506-513.
224. Alexopoulos GS, Raue PJ, Kiosses DN, et al. Problem-solving therapy and supportive therapy in older adults with major depression and executive dysfunction: Effect on disability. *Arch Gen Psychiatry*. 2011;68(1):33-41.

225. Collard RM, Comijs HC, Naarding P, Oude Voshaar RC. Physical frailty: Vulnerability of patients suffering from late-life depression. *Aging Ment Health*. 2014;18(5):570-578.
226. Peerenboom L, Collard RM, Naarding P, Comijs HC. The association between depression and emotional and social loneliness in older persons and the influence of social support, cognitive functioning and personality: A cross-sectional study. *J Affect Disord*. 2015;182:26-31.
227. Shah SN, Meeks S. Late-life bereavement and complicated grief: A proposed comprehensive framework. *Aging Ment Health*. 2012;16(1):39-56.
228. Benraad CE, Kamerman-Celie F, van Munster BC, Oude Voshaar RC, Spijker J, Olde Rikkert MG. Geriatric characteristics in randomised controlled trials on antidepressant drugs for older adults: A systematic review. *Int J Geriatr Psychiatry*. 2016;31(9):990-1003.
229. Gunn J, Diggins J, Hegarty K, Blashki G. A systematic review of complex system interventions designed to increase recovery from depression in primary care. *BMC Health Serv Res*. 2006;6:88-6963-6-88.
230. Bruijniks SJE, Franx G, Huibers MJH. The implementation and adherence to evidence-based protocols for psychotherapy for depression: The perspective of therapists in dutch specialized mental healthcare. *BMC Psychiatry*. 2018;18(1):190-018-1768-x.
231. Sinnema H, Terluin B, Wensing M, et al. Systematic tailoring for the implementation of guideline recommendations for anxiety and depressive disorders in general practice: Perceived usefulness of tailored interventions. *BMC Fam Pract*. 2013;14:94-2296-14-94.
232. Sinnema H, Majo MC, Volker D, et al. Effectiveness of a tailored implementation programme to improve recognition, diagnosis and treatment of anxiety and depression in general practice: A cluster randomised controlled trial. *Implement Sci*. 2015;10:33-015-0210-8.
233. Frost R, Bauernfreund Y, Walters K. Non-pharmacological interventions for depression/anxiety in older adults with physical comorbidities affecting functioning: Systematic review and meta-analysis. *Int Psychogeriatr*. 2019;31(8):1121-1136.
234. Kirkham JG, Choi N, Seitz DP. Meta-analysis of problem solving therapy for the treatment of major depressive disorder in older adults. *Int J Geriatr Psychiatry*. 2016;31(5):526-535.
235. Archer J, Bower P, Gilbody S, et al. Collaborative care for depression and anxiety problems. *Cochrane Database Syst Rev*. 2012;10:CD006525.
236. Beyer JL, Johnson KG. Advances in pharmacotherapy of late-life depression. *Curr Psychiatry Rep*. 2018;20(5):34-018-0899-6.
237. Cuijpers P, Noma H, Karyotaki E, Vinkers CH, Cipriani A, Furukawa TA. A network meta-analysis of the effects of psychotherapies, pharmacotherapies and their combination in the treatment of adult depression. *World Psychiatry*. 2020;19(1):92-107.
238. Huang A, Delucchi K, Dunn LB, Nelson JC. A systematic review and meta-analysis of psychotherapy for late-life depression. *Am J Geriatr Psychiatry*. 2015;23(3):261-273.
239. Markle-Reid M, Browne G, Gafni A. Nurse-led health promotion interventions improve quality of life in frail older home care clients: Lessons learned from three randomized trials in ontario, canada. *J Eval Clin Pract*. 2013;19(1):118-131.
240. Bingham KS, Kumar S, Dawson DR, Mulsant BH, Flint AJ. A systematic review of the measurement of function in late-life depression. *Am J Geriatr Psychiatry*. 2018;26(1):54-72.
241. Rogers JC, Holm MB, Raina KD, et al. Disability in late-life major depression: Patterns of self-reported task abilities, task habits, and observed task performance. *Psychiatry Res*. 2010;178(3):475-479.

242. Karp JF, Skidmore E, Lotz M, Lenze E, Dew MA, Reynolds CF, 3rd. Use of the late-life function and disability instrument to assess disability in major depression. *J Am Geriatr Soc*. 2009;57(9):1612-1619.
243. Higgins JPT, Thomas J, Chandler J, Cumpston M, Page MJ, Welch VA. Cochrane handbook for systematic reviews of interventions version 6.0. [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook). Updated 2019.
244. Craig P, Dieppe P, Macintyre S, et al. Developing and evaluating complex interventions: The new medical research council guidance. *BMJ*. 2008;337:a1655.
245. Sheehan DV, Harnett-Sheehan K, Raj BA. The measurement of disability. *Int Clin Psychopharmacol*. 1996;11 Suppl 3:89-95.
246. Ettinger WH, Jr, Burns R, Messier SP, et al. A randomized trial comparing aerobic exercise and resistance exercise with a health education program in older adults with knee osteoarthritis. the fitness arthritis and seniors trial (FAST). *JAMA*. 1997;277(1):25-31.
247. Von Korff M, Ustun TB, Ormel J, Kaplan I, Simon GE. Self-report disability in an international primary care study of psychological illness. *J Clin Epidemiol*. 1996;49(3):297-303.
248. Chekroud SR, Gueorguieva R, Zheutlin AB, et al. Association between physical exercise and mental health in 1.2 million individuals in the USA between 2011 and 2015: A cross-sectional study. *Lancet Psychiatry*. 2018;5(9):739-746.
249. Cunningham C, O'Sullivan R, Caserotti P, Tully MA. Consequences of physical inactivity in older adults: A systematic review of reviews and meta-analyses. *Scand J Med Sci Sports*. 2020;30(5):816-827.
250. Kim SY, Park JH, Lee MY, Oh KS, Shin DW, Shin YC. Physical activity and the prevention of depression: A cohort study. *Gen Hosp Psychiatry*. 2019;60:90-97.
251. Harvey SB, Overland S, Hatch SL, Wessely S, Mykletun A, Hotopf M. Exercise and the prevention of depression: Results of the HUNT cohort study. *Am J Psychiatry*. 2018;175(1):28-36.
252. Kilil-Drori S, Kilil-Drori AJ, Pira S, Rej S. Exercise intervention for late-life depression: A meta-analysis. *J Clin Psychiatry*. 2020;81(1):10.4088/JCP.19r12877.
253. Rhyner KT, Watts A. Exercise and depressive symptoms in older adults: A systematic meta-analytic review. *J Aging Phys Act*. 2016;24(2):234-246.
254. Hall SE, Lawal OA, Clark CM, et al. Novel approach to characterize heterogeneity in an aerobic exercise intervention. *Med Sci Sports Exerc*. 2019;51(7):1506-1516.
255. Guico-Pabia CJ, Fayyad RS, Soares CN. Assessing the relationship between functional impairment/recovery and depression severity: A pooled analysis. *Int Clin Psychopharmacol*. 2012;27(1):1-7.
256. Hegeman JM, van Fenema EM, Comijs HC, Kok RM, van der Mast RC, de Waal MWM. Effect of chronic somatic diseases on the course of late-life depression. *Int J Geriatr Psychiatry*. 2017;32(7):779-787.
257. Choi SM, Yoon GJ, Jung HJ, Kim BC. Analysis of characteristics affecting instrumental activities of daily living in parkinson's disease patients without dementia. *Neurol Sci*. 2019;40(7):1403-1408.
258. Yates JA, Clare L, Woods RT, Cognitive Function and Ageing Study: Wales. What is the relationship between health, mood, and mild cognitive impairment? *J Alzheimers Dis*. 2017;55(3):1183-1193.

259. Shimbo D, Chaplin W, Crossman D, Haas D, Davidson KW. Role of depression and inflammation in incident coronary heart disease events. *Am J Cardiol*. 2005;96(7):1016-1021.
260. Bruce ML, Sirey JA. Integrated care for depression in older primary care patients. *Can J Psychiatry*. 2018;63(7):439-446.
261. Richards DA, Borglin G. Complex interventions and nursing: Looking through a new lens at nursing research. *Int J Nurs Stud*. 2011;48(5):531-533.
262. Krahn DD, Bartels SJ, Coakley E, et al. PRISM-E: Comparison of integrated care and enhanced specialty referral models in depression outcomes. *Psychiatr Serv*. 2006;57(7):946-953.
263. Wade DT, Halligan PW. The biopsychosocial model of illness: A model whose time has come. *Clin Rehabil*. 2017;31(8):995-1004.
264. Jeste DV, Palmer BW, Rettew DC, Boardman S. Positive psychiatry: Its time has come. *J Clin Psychiatry*. 2015;76(6):675-683.
265. Stoner CR. Positive psychiatry/psychology for older adults: A new and important movement but robust methodology is essential. *Int Psychogeriatr*. 2019;31(2):163-165.
266. Valentijn P. Rainbow of chaos: A study into the theory and practice of integrated primary care. *international journal of integrated care*. 2016;16(2):3.
267. Valentijn P, Opheij W, Bruijnzeels MA. Understanding integrated care: A comprehensive conceptual framework based on the integrative functions of primary care. *Int J Integr Care*. 2013;13:10.
268. Ryan DJ, Willems JA, Stebbings GK, Morse CI, Stewart CE, Onambele-Pearson GL. Reliability and validity of the international physical activity questionnaire compared to calibrated accelerometer cut-off points in the quantification of sedentary behaviour and physical activity in older adults. *PLoS One*. 2018;13(4):e0195712.
269. Dyrstad SM, Hansen BH, Holme IM, Anderssen SA. Comparison of self-reported versus accelerometer-measured physical activity. *Med Sci Sports Exerc*. 2014;46(1):99-106.
270. Kowalski K, Rhodes R, Naylor PJ, Tuokko H, MacDonald S. Direct and indirect measurement of physical activity in older adults: A systematic review of the literature. . 2012;9:148.
271. LeMoult J, Gotlib IH. Depression: A cognitive perspective. *Clin Psychol Rev*. 2019;69:51-66.
272. Schuch FB, Deslandes AC, Stubbs B, Gosmann NP, Silva CTBD, Fleck MPA. Factors that influence the neurobiological effects of exercise likely extend beyond age and intensity in people with major depression. *Neurosci Biobehav Rev*. 2017;77:301-302.
273. Dowd KP, Szeklicki R, Minetto MA, et al. A systematic literature review of reviews on techniques for physical activity measurement in adults: A DEDIPAC study. *Int J Behav Nutr Phys Act*. 2018;15(1):15-017-0636-2.
274. Ainsworth B, Cahalin L, Buman M, Ross R. The current state of physical activity assessment tools. *Prog Cardiovasc Dis*. 2015;57(4):387-395.
275. Gold LH. DSM-5 and the assessment of functioning: The world health organization disability assessment schedule 2.0 (WHODAS 2.0). *J Am Acad Psychiatry Law*. 2014;42(2):173-181.
276. Piek E, van der Meer K, Hoogendijk WJ, Penninx BW, Nolen WA. Most antidepressant use in primary care is justified; results of the Netherlands study of depression and anxiety. *PLoS One*. 2011;6(3):e14784.

277. Weber M, Belala N, Clemson L, et al. Feasibility and effectiveness of intervention programmes integrating functional exercise into daily life of older adults: A systematic review. *Gerontology*. 2018;64(2):172-187.
278. Boulton E, Hawley-Hague H, French DP, et al. Implementing behaviour change theory and techniques to increase physical activity and prevent functional decline among adults aged 61-70: The PreventIT project. *Prog Cardiovasc Dis*. 2019;62(2):147-156.
279. Stallinga HA. Functioneren als kern van verpleegkunde. *TVZ verpleegkunde in praktijk en wetenschap*. 2019;4:32.
280. Kearney PM, Pryor J. The international classification of functioning, disability and health (ICF) and nursing. *J Adv Nurs*. 2004;46(2):162-170.
281. Stallinga HA, Roodbol PF, Annema C, Jansen GJ, Wynia K. Functioning assessment vs. conventional medical assessment: A comparative study on health professionals' clinical decision-making and the fit with patient's own perspective of health. *J Clin Nurs*. 2014;23(7-8):1044-1054.
282. Boonstra N, Roorda S, Aardema H. Werken met de international classification of functioning. *Nurse Academy GGZ*. 2017;1:10.



